

2024-04-04 10:00




FIG. 1






FIG. 2

+








FIG. 5




FIG. 6





FIG. 7








FIG. 9 +




FIG. 10




FIG. 11

+




FIG. 12

10044944.0001




FIG. 13





FIG. 14



FIG. 16






FIG. 17






FIG. 18




FIG. 19




FIG. 20




FIG. 21




FIG. 22




FIG. 23




FIG. 24




FIG. 25



1003494-10394




FIG. 26




FIG. 27




FIG. 28




FIG. 29








FIG. 32


[illegible]


FIG. 33

+

+

91








FIG. 35





FIG. 36

FIG. 37






FIG. 38




FIG. 38 A

FIG. 38B

FIG. 38C

FIG. 38D






FIG. 39



+

FIG. 40



10044944 10004

+





FIG. 41






FIG. 42




FIG. 42A




FIG. 43

1004494-102201




FIG. 43A





FIG. 44

FIG. 45








FIG. 47






FIG. 48




FIG. 49




FIG. 50

FIG. 51





FIG. 52



100444-10001




FIG. 53




FIG. 54




FIG. 55






FIG. 56

10044944.0001




FIG. 57




FIG. 58




FIG. 59




FIG. 61




FIG. 62





FIG. 63









FIG. 64

+






FIG. 66




FIG. 68

FIG. 67




FIG. 69





FIG. 70







FIG. 72



FIG. 73






FIG. 74




FIG. 75




FIG. 76




FIG. 77

2628




FIG. 78

FIG. 79







FIG. 81

+








FIG. 83

+




FIG. 84A





FIG. 84B




FIG. 84C

+




FIG. 84D

FIG. 85




FIG. 85

# I. SEASONAL CHART FOR VINEYARD MECHANIZATION ACTIVITIES OF VITIS LABRUSCANA (and other grapes with drooping growth habits) ON SINGLE CURTAIN TRELLIS




FIG. 86

Exact date of each operation will depend on the viticultural region. The exact date can vary from region to region by as much as 3-4 weeks (depending on the cultivar). Therefore, mechanical operation should be based on physiological growth of the vine. Of course, the seasons in the southern hemisphere are opposite.

## II. SEASONAL CHART FOR VINEYARD MECHANIZATION ACTIVITIES OF VITIS LABRUSCANA (and other grapes with drooping growth habits) ON GDC TRELLIS AND GDC-LIKE CANOPY SYSTEMS




FIG. 87

Exact date of each operation will depend on the viticultural region. The exact date can vary from region to region by as much as 3-4 weeks (depending on the cultivar). Therefore, mechanical operation should be based on physiological growth of the vine. Of course, the seasons in the southern hemisphere are opposite.



### III. SEASONAL CHART FOR VINEYARD MECHANIZATION ACTIVITIES ON MINIMAL PRUNED *VITIS LABRUSCANA* (and other grapes with drooping growth habits) ON SINGLE CURTAIN TRELLIS SYSTEMS




FIG. 88

Exact date of each operation will depend on the viticultural region. The exact date can vary from region to region by as much as 3-4 weeks (depending on the cultivar). Therefore, mechanical operation should be based on physiological growth of the vine. Of course, the seasons in the southern hemisphere are opposite.

# IV. SEASONAL CHART FOR VINEYARD MECHANIZATION ACTIVITIES ON MINIMAL PRUNED VITIS LABRUSCANA (and other grapes with drooping growth habits) ON GDC TRELLIS SYSTEMS




FIG. 89

Exact date of each operation will depend on the viticultural region. The exact date can vary from region to region by as much as 3-4 weeks (depending on the cultivar). Therefore, mechanical operation should be based on physiological growth of the vine. Of course, the seasons in the southern hemisphere are opposite.

# V. SEASONAL CHART FOR VINEYARD MECHANIZATION ACTIVITIES OF VITIS VINIFERA AND FRENCH AMERICAN HYBRIDS PRODUCED ON HIGH WIRE SINGLE CURTAIN TRELLISES




FIG. 90

Exact date of each operation will depend on the viticultural region. The exact date can vary from region to region by as much as 3-4 weeks (depending on the cultivar). Therefore, mechanical operation should be based on physiological growth of the vine. Of course, the seasons in the southern hemisphere are opposite.

# VI. SEASONAL CHART FOR VINEYARD MECHANIZATION ACTIVITIES OF VITIS VINIFERA AND FRENCH AMERICAN HYBRIDS PRODUCED ON GDC AND OTHER DIVIDED CANOPY TRELLISES




FIG. 91

Exact date of each operation will depend on the viticultural region. The exact date can vary from region to region by as much as 3-4 weeks (depending on the cultivar). Therefore, mechanical operation should be based on physiological growth of the vine. Of course, the seasons in the southern hemisphere are opposite.

VII. SEASONAL CHART FOR VINEYARD MECHANIZATION ACTIVITIES IN  
MINIMAL PRUNED VITIS VINIFERA AND FRENCH AMERICAN HYBRIDS  
TRAINED TO A HIGH WIRE SINGLE CURTAIN TRELLISING SYSTEM.




FIG. 92

Exact date of each operation will depend on the viticultural region. The exact date can vary from region to region by as much as 3-4 weeks (depending on the cultivar). Therefore, mechanical operation should be based on physiological growth of the vine. Of course, the seasons in the southern hemisphere are opposite.

# VIII. SEASONAL CHART FOR VINEYARD MECHANIZATION ACTIVITIES ON MINIMAL PRUNED VITIS VINIFERA AND FRENCH AMERICAN HYBRIDS ON GDC TRELLIS SYSTEMS

Jan. - (late) March      April      May      June      July      Aug.      Sept.      Oct.-Dec.




FIG. 93

Exact date of each operation will depend on the viticultural region. The exact date can vary from region to region by as much as 3-4 weeks (depending on the cultivar). Therefore, mechanical operation should be based on physiological growth of the vine. Of course, the seasons in the southern hemisphere are opposite.

[illegible]

Exact date of each operation will depend on the viticultural region. The exact date can vary from region to region by as much as 3-4 weeks (depending on the cultivar). Therefore, mechanical operation should be based on physiological growth of the vine. Of course, the seasons in the southern hemisphere are opposite.

# X. SEASONAL CHART FOR VINEYARD MECHANIZATION ACTIVITIES OF VITIS VINIFERA AND FRENCH AMERICAN HYBRIDS PRODUCED ON STANDARD VERTICAL MOVEABLE CATCH WIRES




FIG. 95

Exact date of each operation will depend on the viticultural region. The exact date can vary from region to region by as much as 3-4 weeks (depending on the cultivar). Therefore, mechanical operation should be based on physiological growth of the vine. Of course, the seasons in the southern hemisphere are opposite.



# XI. SEASONAL CHART FOR VINEYARD MECHANIZATION ACTIVITIES OF VITIS VINIFERA AND FRENCH AMERICAN HYBRIDS PRODUCED ON LYRE OR "U" AND OTHER DIVIDED CANOPY TRELLISES





FIG. 96

Exact date of each operation will depend on the viticultural region. The exact date can vary from region to region by as much as 3-4 weeks (depending on the cultivar). Therefore, mechanical operation should be based on physiological growth of the vine. Of course, the seasons in the southern hemisphere are opposite.

## XII. SEASONAL CHART FOR VINEYARD MECHANIZATION ACTIVITIES OF VITIS VINIFERA AND FRENCH AMERICAN HYBRIDS ON SMART-DYSON BALLERINA (and similar) TRELLISING SYSTEMS.



**FIG. 97**

\*Use on the upper part of Smart-Dyson Ballerina.

\*\*Use on the lower part of Smart-Dyson Ballerina.

Exact date of each operation will depend on the viticultural region. The exact date can vary from region to region by as much as 3-4 weeks (depending on the cultivar). Therefore, mechanical operation should be based on physiological growth of the vine. Of course, the seasons in the southern hemisphere are opposite.